


Long-Term Safety and Efficacy of Oral Deucricitbant for Treatment of Hereditary Angioedema Attacks: Results of the RAPiDe-2 Extension Study

Michael E. Manning¹, John Anderson², Emel Aygören-Pürsün³, Laurence Bouillet⁴, Hugo Chapdelaine⁵, Henriette Farkas⁶, Delphine Gobert⁷, Roman Hakl⁸, Joshua S. Jacobs⁹, Ramon Lleonart¹⁰, Avner Reshef¹¹, Giuseppe Spadaro¹², Maria Staevska¹³, Marcin Stobiecki¹⁴, Anna Valerieva¹³, Justin Sun¹⁵, Li Zhu¹⁵, Ming Yu¹⁵, Giorgio Giannattasio¹⁶, Peng Lu¹⁵

¹Allergy, Asthma and Immunology Associates, Ltd., Scottsdale, AZ, USA; ²AllerVieHealth, Clinical Research Center of Alabama, Birmingham, AL, USA; ³University Hospital Frankfurt, Department for Children and Adolescents, Goethe University Frankfurt, Frankfurt, Germany; ⁴National Reference Center for Angioedema (CREAK), Department of Internal Medicine, Grenoble Alpes University, Laboratoire T-RAIG, UMR 5525 TIMC-IMAG (UGA-CNRS), Grenoble, France; ⁵CHU de Montréal, Université de Montréal, Montréal, QC, Canada; ⁶Hungarian Angioedema Center of Reference and Excellence, Department of Internal Medicine and Haematology, Semmelweis University, Budapest, Hungary; ⁷Sorbonne Université, Médecine Interne, AP-HP, Centre de référence des angioédèmes à kinines, Hôpital Saint-Antoine, Paris, France; ⁸St. Anne's University Hospital in Brno and Faculty of Medicine Department of Clinical Immunology and Allergology, Masaryk University, Brno, Czech Republic; ⁹Allergy and Asthma Clinical Research, Walnut Creek, CA, USA; ¹⁰Bellvitge University Hospital, Allergy Service, L'Hospitalet de Llobregat, Barcelona, Spain; ¹¹Barzilai University Hospital, Allergy, Immunology and Angioedema Center, Ashkelon, Israel; ¹²University of Naples Federico II, Department of Translational Medical Sciences and Center for Basic and Clinical Immunology Research (CISI), Napoli, Italy; ¹³Medical University of Sofia, Department of Allergology, Sofia, Bulgaria; ¹⁴Jagiellonian University Medical College, Department of Clinical and Environmental Allergology, Krakow, Poland; ¹⁵Pharvaris Inc., Lexington, MA, USA; ¹⁶Pharvaris GmbH, Zug, Switzerland

Key takeaways

The ongoing Phase 2/3 RAPiDe-2 extension study provides evidence on the long-term safety and efficacy of deucricitbant immediate-release (IR) capsule for on-demand treatment of repeat hereditary angioedema (HAE) attacks.



Background

- Hereditary angioedema (HAE):** a bradykinin-mediated condition with painful swelling attacks affecting multiple locations in the body.¹
- Current landscape:** guidelines recommend HAE attacks are treated as early as possible.²⁻⁴ Parenteral administration often leads to on-demand treatment of HAE attacks being delayed or forgone.⁵⁻¹⁴
- Oral deucricitbant:** a selective, bradykinin B2 receptor antagonist under development for both prophylactic and on-demand treatment of HAE attacks.¹⁵⁻²²


Objective

Evaluate the long-term safety and efficacy of deucricitbant IR capsule for on-demand treatment of repeat HAE attacks in the RAPiDe-2 extension study.

Methods

- RAPiDe-2 (NCT05396105)*:** a two-part, double-blind Phase 2/3 extension study.¹⁷
- Part A eligible participants:** adults who completed RAPiDe-1 (NCT04618211).¹⁵
- Part A prophylaxis:** no long-term HAE prophylaxis treatment is allowed. Recent use of long-term HAE prophylaxis treatment prior to screening is allowed provided a pre-specified washout period is observed.


Figure 1. RAPiDe-2 study design



Methods

Primary endpoint: Safety including treatment-emergent adverse events (TEAEs), clinical laboratory tests, vital signs, and electrocardiogram findings.

Secondary endpoints: Efficacy endpoints using two patient-reported outcome tools.



- Data collection pre-specified at pre-treatment, hourly for 6 hours, and at 8, 12, 24, and 48 hours post-treatment.

Table 1. Efficacy endpoints

Key efficacy endpoints	Defined as
Time to	
Onset of symptom relief	By 12 hours, PGI-C rating of at least "a little better" for 2 consecutive timepoints ^a
Reduction in attack severity	By 12 hours, ≥1-level reduction in the PGI-S from pre-treatment for 2 consecutive timepoints ^a
Substantial symptom relief	By 12 hours, PGI-C rating of at least "better" for 2 consecutive timepoints ^a
Proportion of attacks achieving	
Complete attack resolution	By 24 hours, PGI-S rating of "none" ^b

PGI-C, Patient Global Impression of Change; PGI-S, Patient Global Impression of Severity. ^aIf rescue medication used within 14.5 hours post-treatment, time to event was censored at 14.5 hours regardless of whether event occurred within 12 hours post-treatment. ^bRescue medication use within 33.5 hours post-treatment was regarded as not achieving complete attack resolution at 24 hours.

Post hoc analyses: Safety and efficacy for on-demand treatment of upper airway attacks, including laryngeal attacks without breathing difficulties.

- Upper airway attacks confirmed by investigators as per protocol definition: Swelling of the lips/tongue or any sensation of lump in the throat, difficulty swallowing, or voice change.
- Difficulty swallowing and voice change were assessed using the 5-symptom composite Angioedema symptom Rating scAle (AMRA-5).

Results

Data:

- Data snapshot from RAPiDe-2 Part A. Combined dose-blinded group data shown.

Safety analysis

- Participants who received ≥1 dose of deucricitbant IR capsule in the study at data cutoff (10 June 2024).
- 337 attacks from 19 participants.

Table 2. TEAEs within 5 days of study drug administration

Adverse events	Deucricitbant IR capsule (Combined dose group)
Attacks with any TEAE, n (%)	13 (3.9)
Treatment-related TEAEs, n	0
Serious TEAEs, n	1 ^a
Treatment-related serious TEAEs, n	0
TEAEs leading to study drug discontinuation, study withdrawal, or death, n	0

IR, immediate-release; TEAE, treatment-emergent adverse event (defined as adverse event occurring during time window from first study drug administration). ^aTooth caries unrelated to treatment.

Results

Efficacy analysis

- Modified intention-to-treat analysis set: participants who treated ≥1 attack with deucricitbant IR capsule and non-missing PGI-C results from ≥1 post-treatment timepoint at data cutoff (01 March 2024).
- 265 attacks from 17 participants.

Figure 3. Median time to achieving key efficacy endpoints



Figure 4. Majority of attacks achieved key efficacy endpoints within timeframe




Figure 5. Over 90% of attacks achieving complete attack resolution within 24 hours were treated with a single dose of deucricitbant





Figure 6. 98.5% of attacks treated with deucricitbant did not use rescue medication



Results

Upper airway attacks analysis

- 337 attacks at data cutoff (31 May 2024), of which 7 were upper airway, including laryngeal, attacks.
 - Difficulty in swallowing and/or voice change were reported as symptom manifestations of 3 attacks before treatment.[†]

Table 3. Similar times to symptom relief for upper airway and other attacks

	Upper airway attacks	Non-upper airway attacks in participants with upper airway attacks	Total non-upper airway attacks
Number of participants	5	5	19
Total number of attacks treated^a	7	116	328
Time to onset of symptom relief^{b,c}			
Number of attacks ^d	7	112	318
Median hours (95% CI)	0.9 (0.5, 2.0)	1.0 (1.0, 1.1)	1.1 (1.0, 1.1)
Time to reduction in attack severity^{e,g}			
Number of attacks ^f	6	111	312
Median hours (95% CI)	3.0 (0.9, NE)	2.0 (2.0, 2.7)	2.7 (2.1, 2.9)

CI, confidence interval; NE, not estimable (insufficient data to calculate reliable estimate); PGI-C, Patient Global Impression of Change; PGI-S, Patient Global Impression of Severity. ^a337 attacks treated by 19 participants at data cutoff (31 May 2024). ^bPGI-C rating of at least "a little better" for 2 consecutive timepoints by 12 hours post-treatment. ^cWithin-participant correlation not accounted for in all Kaplan-Meier estimates. ^dEvaluable attacks include deucricitbant-treated attacks with ≥1 post-treatment PGI-C assessment. ^e≥1-level reduction in PGI-S from pre-treatment for 2 consecutive timepoints by 12 hours. ^fEvaluable attacks included deucricitbant-treated attacks with a pre- and ≥1 post-treatment PGI-S assessment.

Figure 7. Majority of upper airway attacks were treated with a single dose of deucricitbant





Figure 8. Majority of upper airway and other attacks did not use rescue medication by 48 hours post-treatment



References

- Busse PJ, et al. *N Engl J Med.* 2020;382:1136-48.
- Betschel S, et al. *Allergy Asthma Clin Immunol.* 2019;15:72.
- Busse PJ, et al. *J Allergy Clin Immunol Pract.* 2021;9:132-50.
- Maurer M, et al. *Allergy.* 2022;77:1961-90.
- Berliner R. [package insert]. <https://labeling.cslbehring.com/pi/us/berliner/en/berliner-prescribing-information.pdf>. Accessed January 23, 2025.
- Cinzyze. [summary of product characteristics]. https://www.ema.europa.eu/en/documents/product-information/cinzyze-epar-product-information_en.pdf. Accessed January 23, 2025.
- Firazy®. [package insert]. https://www.shirecontent.com/PDFs/Firazy_USA_ENG.pdf. Accessed January 23, 2025.
- Kalbitor®. [package insert]. https://www.shirecontent.com/PDFs/Kalbitor_USA_ENG.pdf. Accessed January 23, 2025.
- Ruconest®. [package insert]. https://www.ruconest.com/wp-content/uploads/Ruconest_PL_Apr2020.pdf. Accessed January 23, 2025.
- Burnette A, et al. Presented at: AAAAI; February 24-27, 2023; San Antonio, TX, USA.
- Betschel SD, et al. *Allergy Asthma Clin Immunol.* 2024;20:43.
- Center for Biologics Evaluation and Research. The voice of the patient—hereditary angioedema. US Food and Drug Administration; May 2018. <https://www.fda.gov/media/113509/download>. Accessed January 23, 2025.
- Radiojicic C, et al. Presented at: AAAAI; February 24-27, 2023; San Antonio, TX, USA.
- Mendivil J, et al. Presented at: ACAAI; November 9-13, 2023; Anaheim, CA, USA.
- APiDe-1. <https://www.clinicaltrials.gov/study/NCT04618211>. Accessed January 23, 2025.
- Maurer M, et al. Presented at: AAAAI; February 24-27, 2023; San Antonio, TX, USA.
- RAPiDe-2. <https://clinicaltrials.gov/study/NCT05396105>. Accessed January 23, 2025.
- RAPiDe-3. <https://www.clinicaltrials.gov/study/NCT06343779>. Accessed January 23, 2025.
- CHAPTER-1. <https://www.clinicaltrials.gov/study/NCT05074185>. Accessed January 23, 2025.
- CHAPTER-3. <https://clinicaltrials.gov/study/NCT06669754>. Accessed January 23, 2025.
- CHAPTER-4. <https://www.clinicaltrials.gov/study/NCT06679881>. Accessed January 23, 2025.
- Aygören-Pürsün E, et al. Presented at: EAACI; May 31-June 3, 2024; Valencia, Spain.
- Cohn DM, et al. *Clin Transl Allergy.* 2023;12:288.

This presentation includes data for an investigational product not yet approved by regulatory authorities.